



Indonesian Food Science and Technology Journal

**INDONESIAN FOOD SCIENCE AND
TECHNOLOGY JOURNAL (IFSTJ)**Journal homepage: online-journal.unja.ac.id/ifstj/issue/archive

The Technological Research Outcomes of Boiled Sausages Fortified with Sea Buckthorn Peels

Erkigul Bukyei^{1#}, Buyanchimeg Baasanjargal¹, Enkhbat Tumurbaatar¹, Altanzul Ragchaa¹,
Solongo Ganbold²

¹ School of Engineering and Technology, Mongolian University of Life Sciences, Zaisan, Ulaanbaatar, 17024, Mongolia.

² Mongolian University of Science and Technology, Baga Toiruu 34, Ulaanbaatar, 14191, Mongolia

#Corresponding author: E-mail: erkigul@mul.edu.mn

Abstract — The article presents the evaluation results on dried peels from sea buckthorn fruit pomace obtained after juice production in Tes Sum, Uvs province, and its use in producing boiled sausages. The boiled sausages were enriched with four different amounts of dried sea buckthorn peels, which are as follows: 0.2%, 0.3%, 0.4%, and 0.5%. The fortified sausages had been evaluated for chemical composition. According to the experimental research, compared to the reference, the amount of oil increased by 1.9%, fiber by 0.4%, flavonoids by 0.2mg%, calcium and magnesium by 2mg%, and iron by 0.2% in boiled sausages enriched with 0.4% sea buckthorn peels. Moreover, the moisture content and nitrite levels of fortified sausages with sea buckthorn peels decreased by 3.3% and 2.2mg/ % respectively.

Keywords — sea buckthorn peels, meat products, fortifier, technology

Manuscript received Oct 04, 2022; revised Feb 20, 2023; accepted June 10, 2023. Available online July 30, 2023.

Indonesian Food Science and Technology Journal is licensed under a Creative Commons Attribution 4.0 International License



I. INTRODUCTION

Within the framework of the "Sea buckthorn" program implemented by the Government of Mongolia, projects include increasing fruit cultivation, improving harvesting technology, refining industrial processing technology, increasing the types of products that are produced from sea buckthorn fruit, and conducting clinical studies on the biochemical composition of the product and its effects on human health have been developed and carried out through the collaboration of enterprises, organizations, and academic centers [1].

Although the result of the program reflects the increase in enterprises that plant sea buckthorn and produce products in Mongolia, these enterprises only separate the juice and oil mechanically and the soft tissue and seeds and peels are wasted or used as animal feed. Sea buckthorn pomace, which is considered as waste matter, contains a significant amount of biologically active compounds. There is a focus on processing and using it as a fortification in production [2].

In our country, imported peas, red and green sweet peppers mainly play the part of fortifier in the production of sausages

and meat products. There is a reason to believe that almost no variety of sausages with plant additions, except for the boiled sausages called "Pepper" and "Pea Sausage," which are produced by German technology and supplied to the market.

Enriching food items with substances that are extremely important to the human body should be treated carefully and not by common principles, but by taking into account several factors such as the conditions of the country, the customs, and traditions of the people, the nature and climate [3].

Therefore, in the research work, an experimental study was conducted to develop boiled sausage technology enriched with sea buckthorn peels grown in an ecologically clean environment that has preserved its natural appearance and increased the biological value of sausages.

The aim of the research is to develop a new technology that improves the structure and quality of boiled sausages by using dried sea buckthorn peels in the sausage production process. To achieve this, we set the following goals. It includes: study the chemical composition of dried sea buckthorn peel, optimize the amount of sea buckthorn peel added into making boiled sausages and determine and evaluate the physical and chemical parameters of enriched-sausages.

II. MATERIAL AND METHODS

A. Materials

Dried Sea buckthorn peels from the fruit pomace obtained after juice production in Tes Sum, Uvs province, have been used to prepare the sausages.

B. Methods

The sausage has been made according to the procedure depicted in Figure 1.




Fig. 1 Flow chart for sausage processing

Determinations of moisture, fat, protein, and ash contents in the sausage samples were performed according to MNS ISO 6477:2014, 1444:1984, 937:1984, and 9376:2003, respectively. The microbiological analyses were performed following the MNS 6308:2012, MNS ISO 6579-1:2020, 4833:1995 as well as 6579:1995. The analysis of the mineral composition of sausages was performed by atom spectrophotometry. The crude fiber of sausage was determined according to AOAC 978.10.

The rotatable central composite design was used in the experiment design and the obtained experimental data were processed using the STATISTICA 12 software package.

The research was carried out at the Food Research Laboratory of Mongolian University of Life Science (MULS), the Food biochemistry laboratory of the Institute of Technology, and the accredited food testing laboratory of the "CAMO" Institute using standard methods of instrumental analysis. The sausage-making experiments were performed at the meat processing center and plant of the MULS and Khan Brand LLC. The sensory properties of the fortified sausages (appearance, color, and flavor) were estimated using a quantitative-descriptive test with a grading scale from 2 to 14 by 28 untrained panelist.

III. RESULT AND DISCUSSION

A. Characterization of sea buckthorn peels

Sea buckthorn peels were separated from the fruit pomace and were dried in a drying chamber until their moisture content reached 6% to evaluate their chemical composition. Table 1 depicts the chemical composition of sea buckthorn peels.

TABLE 1
CHEMICAL CHARACTERISTICS OF DRIED SEA BUCKTHORN PEELS, %

No	Chemical characteristics	Dried sea buckthorn peels	G. Nanjid [4]	M.N. Volgarev [6]
1	Moisture	6.2±0.2	6.2	5.8
2	Lipid	30.0±0.4	15.0	8
3	Protein	24.4±0.3	25.49	26
4	Fiber	14.0±0.3	12.3	24
5	Ash	1.7±0.2	0.74	2.9

The data obtained from the experimental studies corresponded with those previously reported results of G. Nanjid et al. [4], [5] and M.N. Volgarev et al. [6].

From the experimental data presented in Table 1, it is apparent that the fiber, ash, moisture, and protein content of the dried sea buckthorn peels are comparable to the results of G. Nanjid et al. [4]. However, fiber content and ash levels were less than 10% and 1.2%, respectively, than the results of M.N. Volgarev et al. [6], which directly relates to its oil content.

In addition, the oil content of the peels was 2 and 4 times higher than the result of previous work [4], [6]. This discrepant result may be due to the oil separation technology. In particular, there is a lack of efficient methods for separating oil from sea buckthorn peels in domestic industry.

The mineral contents and the total flavonoid content in sea buckthorn peels are in Table 2.

TABLE 2
THE CONTENT OF SOME MINERALS AND FLAVONOIDS OF DRIED SEA BUCKTHORN PEELS, mg/ %

Minerals	Dried sea buckthorn peels	Dried sea buckthorn peels [6]	Dried apple [6]	Dried raisins [6]
Calcium	21.0±0.6	22±1.0	11.0±0.6	8.0±0.8
Magnesium	11.0±0.05	9.0±1.5	7.0±1	10.0±1.5
Iron	6.6±0.2	7.0±0.5	6.0±0.9	3.0±1
Chloride	34.5±0.8	32.0±1	47.0±0.7	59.0±0.5
Flavonoids	4.1±0.3	ND	ND	ND

According to the results, the amount of calcium, magnesium, and iron chloride contents of the sausage were comparable with that of the sausages reported by M.N. Volgarev et al. [6]. However, no significant differences have been observed in the magnesium, iron, and chloride contents of the sausage. The calcium content was twice higher than in the other dried fruit peels [6]. Furthermore, in the experimental dried peel, 4.1-4.4 mg/% flavonoids were determined.

Based on the results of this research, sea buckthorn peels are highly valued for their essential nutrients for the human body.

B. Characterization of the boiled sausages fortified with sea buckthorn peels

The preparation of sausage in this study used fortification level treatment of sea buckthorn fruit peels: 0.2%, 0.3%, 0.4%, and 0.5% per 100g of meat. The technological processing of the boiled sausages took place in compliance with general technical requirements of sausage products, and nitrite levels in the prepared boiled sausages were two times lower than in the control samples.

On the one hand, the quality of sausages directly depends on consumer appreciation. On the other hand, one of the tasks of this work is finding optimal sea buckthorn peel fortification of the boiled sausages. Therefore, sensory analysis was performed by 28 untrained panelists, including lecturers and

students. The results of sensory tests by panelists on boiled sausage fortified with sea buckthorn peels are presented in Table 3.

TABLE 3
RESULTS OF THE SENSORY EVALUATION OF THE BOILED SAUSAGES FORTIFIED WITH SEA BUCKTHORN PEELS

№	Fortification level, %	Variable			Total score
		Appearance 12-14 score	Colour 2-4 score	Flavor 11-14 score	
1	0.2%	13.2	2.1	11.8	27.1
2	0.3%	13.3	2.9	12.1	28.3
3	0.4%	13.8	3.8	13.5	31.1
4	0.5%	13.6	2.1	12.2	27.9

The study results convey, the sea buckthorn peel fortification level of 0.4%, was given the highest score of 31.1, while the fortification of 0.2%, was given the lowest score of 27.1 by panelists. In addition, eighty percent of all panelists stated that the fortification of boiled sausage through a level of 0.4% met the consumer-consumption characteristics of taste, texture, and color. The boiled sausage enriched with 0.5% of sea buckthorn peels is marked as having a bright yellow color.




Fig. 2 Experimental design

In the technological experiment, input value (peel amount -x₁, nitrate salt-x₂, ice amount-x₃) and output value (fiber-y₁, flavonoid- y₂, nitrate amount- y₃, physical condition-y₄) have been chosen to optimize the peel ingredient. Also, the experimental reliability has been established by a rotatable central composite design of 3 factors (n=20) in numerical data. After that, the curvilinear regression equation was defined.

$$\begin{aligned}
 Y_1 &= 0.58 + 0.159 * x_1 + 0.045 * x_1^2 + 0.018 * x_2 - 0.131 * x_2^2 - 0.044 * x_3 - 0.165 * x_3^2 + 0.075 * x_1x_2 + 0.025 * x_1x_3 \\
 &\quad - 0.075 * x_2x_3 \\
 Y_2 &= 0.212 + 0.058 * x_1 + 0.103 * x_1^2 - 0.071 * x_2 + 0.148 * x_2^2 + 0.036 * x_3^2 + 0.1 * x_1x_2 \\
 Y_3 &= 2.142 - 0.132 * x_1 - 0.331 * x_1^2 - 0.131 * x_2 + 0.092 * x_2^2 + 0.113 * x_3 + 0.088 * x_3^2 - 0.1 * x_1x_2 - 0.05 * x_1x_3 \\
 &\quad - 0.15 * x_2x_3 \\
 Y_4 &= 5.017 - 0.573 * x_1 - 0.312 * x_1^2 + 0.004 * x_2 - 0.321 * x_2^2 + 0.179 * x_3 + 0.214 * x_3^2 + 0.333 * x_1x_2 - 0.583x_1x_3 \\
 &\quad - 0.083 * x_2x_3
 \end{aligned}$$

The significance of Regression coefficients and Student t-tests have been determined by using program software. Additionally, the Fisher F- test has been used to determine whether the model form matched or not (Table 4).

TABLE 4
 RECIPE OPTIMIZATION OF SAUSAGES WITH SEA BUCKTHORN PEELS
 (CRITERIA FOR REGRESSION MODEL RELIABILITY)

Criteria	Results			
	Y ₁	Y ₂	Y ₃	Y ₄
R	0.907	0.912	0.897	0.879
R ²	0.822	0.833	0.791	0.812
F _{experiment}	3.385	5.551	2.562	4.345
F _{theory(0,05; 9; 10)}	0.035	0.006	0.033	0.008

As can be seen in Table 4, the regression determination coefficient is R²=0.791-0.833. On the other hand, assuming the factors influencing Y₁, Y₂, Y₃, and Y₄ are 100%, the dependence of X₁, X₂, and X₃ on factors is 79-83%, while uncounted factors or external influence is low and around 17-21%.

The main influence factors on the technology of boiled sausage with sea buckthorn peel production recipes are displayed on three-dimensional surfaces (Fig. 3)




Fig. 3 Three-dimensional surface of the main influence factors

The main factors that influence the technology of boiled sausage with sea buckthorn peel were estimated based on the results of the mathematical-statistical analysis. The results reveal that amount of dried sea buckthorn peel (X₁) is 0.42%, nitrate salt (X₂) is 0.87%, the amount of ice (X₃) is 22.3%, sausage fiber (Y₁) is 0.48%, flavonoid (Y₂) is 0.2%, nitrate (Y₃) is 2.12 mg, physical condition (Y₄) is 5 points, respectively.

Based on the research results above, we compared the chemical composition of the sausage enriched with 0.4% sea buckthorn peel with the control sample (without sea buckthorn peels) (Figure 4).




Fig. 4 Chemical composition of boiled sausage enriched with sea buckthorn peel and control sample

Compared to the control sample, the sausage enriched with sea buckthorn peel had a higher fat content; the total increase in fat was (1.9%), and the lower moisture content; the total decrease in moisture was (3.3%). The moisture change increases the fat content, and hence the fiber content of the sea buckthorn peel increases its water-binding capacity. The amount of nitrite compared to the controls' was determined to be 2.2 mg/% lower.

Furthermore, the calcium, magnesium, and iron contents of the enriched sausage with sea buckthorn peels increased by 2 mg/%, 2 mg/%, and 0.2 mg/%, separately, compared to the control sample.

TABLE 5
 RESULT OF A MICROBIOLOGICAL STUDY ON THE ENRICHED SAUSAGE WITH SEA BUCKTHORN PEELS

No	Characteristics	Record	Requirements	Enriched sausage
1	Total number of microorganisms per gram	MNS 6308:2012 ND	2*10 ³	<1*10 ²
2	<i>E.coli</i> per gram	ND	ND	Undetected
3	<i>Salmonella</i> per 25 gram		Undetected	Undetected

It should also be notable that after three days of storage at 10°C, the enriched sausage with the addition of reduced nitrite met the standard requirements for the number of bacteria and some pathogenic microorganisms.

IV. CONCLUSION

Based on the results obtained from the experimental investigation on the technology of boiled sausage with sea buckthorn peel, the following conclusions were made: The sea buckthorn peel sample contains 30% oil, 14% fiber, and 1.7% ash (including iron 6.6 mg/%, calcium 21 mg/%, chloride 34.5 mg/%), and flavonoid 4.1 mg/%, indicating that it is a raw material with high biological value. Based on the results of mathematical modeling of sensory evaluation, chemical composition, and technological test parameters optimization of the enriched boiled sausage, it was determined that it is suitable to enhance the boiled sausages with 0.4% sea buckthorn peel. When the boiled sausage was enriched with 0.4% sea buckthorn peel, the average oil content increased by 1.9%, fiber by 0.4%, flavonoids by 0.2mg/%, calcium and magnesium by 2mg/%, and iron by 0.2mg/%, clear evidence of biological value increase. Moreover, sea buckthorn peels used as a fortifier lead to reduced nitrite by 2.2 m/% and moisture content by an average of 3.3%.

ACKNOWLEDGMENTS

We want to express our gratitude to the project "Improving Sea buckthorn fruit processing technology," which is being implemented by the Ministry of Agriculture and Rural Development with the funding of the "Asian Food and Agriculture Cooperation Initiative" (AFACI) international organization.

REFERENCES

- [1] Ministry of Agriculture and Light Industry. 2017. Report on "Chatsargana" National Program 2010–2016
- [2] Ch. Avdai, "Sea Buckthorn" Ulaanbaatar, 2002
- [3] T. Davaa, N. Chimgee, "Canker-berry enriched sausage technology" research, Ulaanbaatar, 2016
- [4] G. Nanjid, "A study on the technology of obtaining sea buckthorn pomace fortified protein-rich dairy products" Ulaanbaatar, 2013
- [5] D.Nasanjargal, "Sea buckthorn processing technology" hand guide, Ulaanbaatar, 2020
- [6] M.N. Volgarev, Chemical composition of food products, 2016, Mosco